

AS1056 - Mathematics for Actuarial Science. Chapter 17, Tutorial 1.

Emilio Luis Sáenz Guillén

Faculty of Actuarial Science
and Insurance,
Bayes Business School.

02/04/2025



Refreshing some concepts: Second-order linear ODEs

A **second-order linear differential equation** is one of the form:

$$\frac{d^2y(x)}{dx^2} + a(x)\frac{dy}{dx} + b(x)y = f(x)$$

The general solution of a second-order linear ODE can be written as:

$$y(x) = \underbrace{\eta(x)}_{\text{particular integral}} + \underbrace{y_0(x)}_{\text{complementary function}}$$

where η and y_0 respectively satisfy:

$$\rightarrow \eta''(x) + a(x)\eta'(x) + b(x)\eta(x) = f(x)$$

$$\rightarrow y_0''(x) + a(x)y_0'(x) + b(x)y_0(x) = 0$$

How do we find $\eta(x)$ and $y_0(x)$?

- ＊ $\eta(x)$: generally a matter of guesswork, guided by the form of the function f .
- ＊ $y_0(x)$: in case of *constant coefficients*, *auxiliary equation* method.

The **auxiliary equation** method provides a straightforward way to find solutions to the complementary function of a linear ODEs with constant coefficients.

$$\lambda^2 + a\lambda + b = 0$$

which has roots λ_1 and λ_2 .

Depending on the sign of the discriminant we have:

- If $a^2 - 4b > 0$, $\lambda_1 \neq \lambda_2$ and:

$$\rightarrow y_0(x) = Ae^{\lambda_1 x} + Be^{\lambda_2 x}$$

- If $a^2 - 4b < 0$, $\lambda_1 = \mu + i\nu$, $\lambda_2 = \mu - i\nu$ and:

$$\rightarrow y_0(x) = e^{\mu x} (Ae^{i\nu x} + Be^{-i\nu x}) = e^{-2x} (C \cos(\nu x) + D \sin(\nu x))$$

- If $a^2 - 4b = 0$, $\lambda_1 = \lambda_2 = \mu$ and:

$$\rightarrow y_0(x) = (Ax + B) e^{\mu x}$$

Exercise 17.13

An object is taken up to a height 10 km and is dropped. It accelerates under gravity but is subject to air resistance, which is proportional to the square of the speed. Let $x(t)$ represent the height of the object at time t , $v(t)$ its speed in a downwards direction. Then

$$\frac{dx}{dt} = -v(t), \quad \frac{dv}{dt} = g - cv(t)^2$$

- (i) Solve the second equation by partial fractions if $v(0) = 0$.
- (ii) Show that $\lim_{t \rightarrow \infty} v(t) = \sqrt{g/c}$. (This is the concept of “terminal velocity”, a speed when deceleration due to air pressure balances the acceleration due to gravity.)

(iii) Show that $x(t)$, measured in metres, is given by

$$x(t) = 10^4 + \frac{\ln(2)}{c} - \frac{1}{c} \ln \left(e^{t\sqrt{gc}} + e^{-t\sqrt{gc}} \right)$$

(iv) Use the values $g = 10\text{ms}^{-2}$, $c = 0.001\text{m}^{-1}$ to show that the object hits the ground approximately 100 seconds after it was released.

Exercise 17.11

Two functions $x(t)$ and $y(t)$ satisfy the differential equations $\frac{dx}{dt} = 4xy - x$, $\frac{dy}{dt} = 1 + \ln(x)$.

(i) Define $X(t) = \ln(x(t))$. Rewrite the two DEs in terms of X and y .

$$\begin{cases} \frac{dX}{dt} = \frac{1}{x} \frac{dx}{dt} = 4y - 1 \\ \frac{dy}{dt} = 1 + \ln(e^X) = 1 + X \end{cases}$$

(ii) Obtain a second order DE for X .

$$\frac{d^2X}{dt^2} = 4 \frac{dy}{dt} = 4(1 + X) \text{ or } \ddot{X} - 4X = 4$$

(iii) Show that $X(t) = -1$ is a particular integral and use this to find the general solution for X and y .

(iv) Evaluate the arbitrary constants in the case where the boundary conditions are $x(0) = 1$, $y(0) = 1$. Hence write down the solutions $x(t)$ and $y(t)$ for all $t \geq 0$.

Bayes Business School

106 Bunhill Row

London EC1Y 8TZ

Tel +44 (0)20 7040 8600

bayes.city.ac.uk