

AS1056 - Mathematics for Actuarial Science. Chapter 9, Tutorial 2.

Emilio Luis Sáenz Guillén

Faculty of Actuarial Science
and Insurance,
Bayes Business School.

12/12/2024



Refreshing some concepts: Geometric series

$$S_n = \sum_{k=0}^{n-1} ar^k = \sum_{k=1}^n ar^{k-1} = \begin{cases} a \left(\frac{1-r^n}{1-r} \right), & \text{for } r \neq 1 \\ an, & \text{for } r = 1 \end{cases}$$

In particular for $n \rightarrow \infty$ we have that:

$$\begin{aligned} \lim_{n \rightarrow \infty} S_n &= \sum_{k=0}^{\infty} ar^k = \sum_{k=1}^{\infty} ar^{k-1} = \lim_{n \rightarrow \infty} a \left(\frac{1-r^n}{1-r} \right) = \\ &= \frac{a}{1-r}, \text{ for } |r| < 1 \end{aligned}$$

- ✿ A geometric series converges if and only if the absolute value of the common ratio is less than 1, i.e., $|r| < 1$.

Exercise 9.1

What is the radius of convergence, ρ , of the series $S_n = \sum_{i=1}^n (2x^2)^i$?

What is the limit of the series when $|x| < \rho$?

For **geometric series** remember the formula,

$$\sum_{k=0}^n x^k = \frac{1 - x^{n+1}}{1 - x}$$

the other formulas can be derived from it:

$$\sum_{k=1}^n x^k = \frac{1 - x^{n+1}}{1 - x} - 1 = \frac{x - x^{n+1}}{1 - x}$$

$$\sum_{k=m}^n x^k = \sum_{k=0}^n x^k - \sum_{k=0}^{m-1} x^k = \frac{x^m - x^{n+1}}{1 - x}$$

Proof by induction

Mathematical induction is a method for proving that some statement $P(n)$ is true for any natural number n .

A **proof by induction** of some statement $P(n)$ consists of two steps:

- 1. Base Case:** Prove the statement $P(n)$ for an initial value of n , usually, $n = 0$ or $n = 1$. The purpose of this step is to show that the statement holds for the first element in the sequence.
- 2. Induction Step:** Assume that the statement $P(n)$ is true for some arbitrary natural number $n = k$ (induction hypothesis). Finally, prove that, if $P(k)$ is true, then the next case, $P(k + 1)$, also holds.

If the statement is true for one case (the base case) + if we assume that the statement holds for an arbitrary case k implies it holds for the next case $k + 1$:

⇒ Then the statement must be true for the entire sequence, i.e., $\forall n \in \mathbb{N}$.

In simpler terms, think of it like dominoes. The base case is like knocking over the first domino, and the inductive step ensures that if one domino falls, the next one will too. Therefore, all the dominoes (or natural numbers) will fall.



Exercise 9.7

Let $f(x) = (2 + x)^{-2}$.

(i) Show that the n th derivative of f is

$$f^{(n)}(x) = (-1)^n(n + 1)!(2 + x)^{-(n+2)}.$$

Refreshing some concepts: Ratio Test

Proposition 8.1 (Ratio Test)

Given a series $S_n = \sum_{n=1}^n a_n$ and the limit

$$\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = r$$

The ratio test states that, the series S_n ,

- converges absolutely —i.e. $\sum_{n=0}^{\infty} |a_n| = L$, for $L \in \mathbb{R}$ —, if $r < 1$;
- diverges, if $r > 1$;
- if $r = 1$ or the limit fails to exist, then the test is inconclusive, because there exist both convergent and divergent series that satisfy this case.


Exercise 9.7

(ii) Evaluate $f^{(n)}(0)$ for each $n > 0$.

(iii) A series $S_n(x)$ is defined by

$$S_n(x) = \frac{1}{4} + \sum_{i=1}^n a_i x^i, \quad \text{where } a_i = f^{(i)}(0).$$

What is the radius of convergence of the series S_n ?



Bayes Business School

106 Bunhill Row

London EC1Y 8TZ

Tel +44 (0)20 7040 8600

bayes.city.ac.uk